ONDERSTEPOORT JOURNAL OF VETERINARY RESEARCH

Volume 67 • Numbers 1–4 • 2000

Author index

AGNAGNA, M.

Aerobic intestinal flora of wild-caught African dwarf crocodiles *Osteolaemus tetraspis*—F.W. HUCHZERMEYER, M.M. HENTON, J. RILEY and M. AGNAGNA 201–204

AIRE, T.A.

Ultrastructural study of the luminal surface of the ducts of the epididymis of gallinaceous birds—T.A. AIRE and D. JOSLING 191–199

AITCHISON, H.

The prevalence of different African horsesickness virus serotypes in the Onderstepoort area near Pretoria, during an outbreak of African horsesickness in South Africa in 1995/1996—C.W. BREMER, G.H. GERDES, H. AITCHISON, I. LOUW, R.R. GREYLING and J. WELGEMOED 65–70

AMANFU, W.

Bovine borreliosis in Botswana—S.P. SHARMA, W. AMANFU and T.C. LOSHO 221–223

ANJILI, C.O.

Leishmania major infections in *Phlebotomus duboscqi* fed on murine models immunized with *L. major* sub-cellular antigens and sandfly gut antigens—P.A. MBATI, C.O. ANJILI, S. ODONGO, P. OGAJA and W. TONU 57–63

ANOSA, V.O.

In vitro erythrophagocytosis by cultured macrophages stimulated with extraneous substances and those isolated from the blood, spleen and bone marrow of Boran and N'Dama cattle infected with *Trypanosoma congolense* and *T. vivax*—V.O. TAIWO and V.O. ANOSA 273–287

Differential expression of surface membrane antigens on bovine monocytes activated with recombinant cytokines and during *Trypanosoma congolense* infection—V.O. TAIWO, V.O. ANOSA and J.O. OLUWANIYI 289–296

BATH, G.F.

The effect of a natural maedi-visna virus infection on the productivity of South African sheep—B. DUNGU, J. VORSTER, G.F. BATH and D.W. VERWOERD 87–96

BENGIS, R.G.

The rise and fall of tuberculosis in a free-ranging chacma baboon troop in the Kruger National Park—D.F. KEET, N.P.J. KRIEK, R.G. BENGIS, D.G. GROBLER and A. MICHEL 115–122

BINTA, M.G.

Helminth parasites of domestic pigeons (*Columba livia domestica*) in Sebele, Gaborone, Botswana—E.Z. MUSHI, M.G. BINTA, R.G. CHABO, R. NDEBELE and R. PANZIRAH 75–76

BLIGHT, G.W.

Evaluation of a commercially available ELISA kit for detection of antibodies to *Anaplasma marginale* and *Anaplasma centrale* in cattle in Australia and Zimbabwe—P.M. BOWLES, J.B. MOLLOY, G.W. BLIGHT, S. SINGH and L.G. MABIKACHECHE 83–86

BOOMKER, J.

A check-list of the nematode parasites of South African serpentes (snakes) and sauria (lizards)—S.F.B.N. HERING-HAGENBECK and J. BOOMKER 1–13

Parasites of South African wildlife. XVI. Helminths of some antelope species from the Eastern and Western Cape Provinces—J. BOOMKER, I.G. HORAK, R. WATERMEYER and D.G. BOOYSE 31–41

Selected chemical parameters in the blood and metals in the organs of the Nile crocodile, *Crocodylus niloticus*, in the Kruger National Park—D. SWANEPOEL, J. BOOMKER and N.P.J. KRIEK 141–148

Molineus cati n. sp. (Nematoda, Trichostrongylina, Molineoidea), a parasite of feral cats, *Felis catus* Linnaeus, 1758 in South Africa—MARIE-CLAUDE DURETTE-DESSET, J. BOOMKER and F.S. MALAN 173–177

Studies on the genus *Setaria* Viborg, 1795 in South Africa. I. *Setaria africana* (Yeh, 1959)—R. WATERMEYER, J. BOOMKER and J.F. PUTTERILL 229–234

BOOYSE, D.G.

Parasites of South African wildlife. XVI. Helminths of some antelope species from the Eastern and Western Cape Provinces—J. BOOMKER, I.G. HORAK, R. WATERMEYER and D.G. BOOYSE 31–41

BOTHA, C.J.

Investigation of biological samples for monofluoroacetate and *Dichapetalum cymosum* poisoning in southern Africa—P.P. MINNAAR, R.I. McCRINDLE, T.W. NAUDÉ and C.J. BOTHA 27–30

BOWLES, J.

Evaluation of a commercially available ELISA kit for detection of antibodies to *Anaplasma marginale* and *Anaplasma centrale* in cattle in Australia and Zimbabwe—P.M. BOWLES, J.B. MOLLOY, G.W. BLIGHT, S. SINGH and L.G. MABIKACHECHE 83–86

BRAACK, L.E.O.

Parasites of domestic and wild animals in South Africa. XXXVIII. Ixodid ticks collected from 23 wild carnivore species—I.G. HORAK, L.E.O. BRAACK, L.J. FOURIE and JANE B. WALKER 239–250

BREMER, C.W.

The prevalence of different African horsesickness virus serotypes in the Onderstepoort area near Pretoria, during an outbreak of African horsesickness in South Africa in 1995/1996—C.W. BREMER, G.H. GERDES, H. AITCHISON, I. LOUW, R.R. GREYLING and J. WELGEMOED 65–70

CHABO, R.G.

Helminth parasites of domestic pigeons (*Columba livia domestica*) in Sebele, Gaborone, Botswana—E.Z. MUSHI, M.G. BINTA, R.G. CHABO, R. NDEBELE and R. PANZIRAH 75–76

CHITAURO, D.

Pasteurella gallinarum: Zimbabwean experience of a versatile pathogen—K. MOHAN, F. DZIVA and D. CHITAURO 301–305

COETZEE, L.

Oribatid mites (Acari, Oribatida) as intermediate hosts of tapeworms of the Family Anoplocephalidae (Cestoda) and the transmission of *Moniezia expansa* cysticercoids in South Africa—R. SCHUSTER, L. COETZEE and J.F. PUTTERILL 49–55

DAKWA, C.

Fertility in rats immunized with steroid-free bovine follicular fluid—C.R. SEWANI-RUSIKE and C. DAKWA 257–262

DICKS, L.M.T.

Phage types of *Salmonella enterica* serovar Enteritidis isolated in South Africa from 1991–1995—L. MARÉ, M.L. VAN DER WALT and L.M.T. DICKS 129–133

DIPPENAAR, S.M.

Observations on a lesser-known monogenean, *Udonella myliobati*, from a copepod parasite, *Lepeophtheirus natalensis*, parasitizing the spotted ragged-tooth shark, *Carcharias taurus*, from South African waters—P.A.S. OLIVIER, S.M. DIPPENAAR, L.F. KHALIL and N.M. MOKGALONG 135–140

DU PREEZ, J.H.

Parameters for the determination and evaluation of heat stress in dairy cattle in South Africa—J.H. DU PREEZ 263–271

DU PLESSIS, L.

The influence of *Mannheimia haemoloytica* A1 seed culture inoculum cell density on the production of leukotoxin in submerged culture supernatant—M.W. ODENDAAL and L. DU PLESSIS 205–216

DUNGU, B.

The effect of a natural maedi-visna virus infection on the productivity of South African sheep—B. DUNGU, J. VORSTER, G.F. BATH and D.W. VERWOERD 87–96

DURETTE-DESSET, MARIE-CLAUDE

Molineus cati n. sp. (Nematoda, Trichostrongylina, Molineoidea), a parasite of feral cats, *Felis catus* Linnaeus, 1758 in South Africa—MARIE-CLAUDE DURETTE-DESSET, J. BOOMKER and F.S. MALAN 173–177

DZIVA, F.

Capsular serogroups of *Pasteurella multocida* isolated from animals in Zimbabwe—F. DZIVA, K. MOHAN and A. PAWANDIWA 225–228

Pasteurella gallinarum: Zimbabwean experience of a versatile pathogen—K. MOHAN, F. DZIVA and D. CHITAURO 301–305

FOURIE, L.J.

Parasites of domestic and wild animals in South Africa. XXXVIII. Ixodid ticks collected from 23 wild carnivore species—I.G. HORAK, L.E.O. BRAACK, L.J. FOURIE and JANE B. WALKER 239–250

GATHUMA, J.M.

Application of ultrasonography in prevalence studies of hydatid cysts in goats in north-western Turkana, Kenya and Toposaland, southern Sudan—E.M. NJOROGE, P.M.F. MBITHI, J.M. GATHUMA, T.M. WACHIRA, J.K. MAGAMBO and E. ZEYHLE 251–255

GATHUMBI, P.K.

Suspected sheep-associated malignant catarrhal fever in a zero-grazed dairy herd in Kenya—C.M MULEI, P.K. GATHUMBI and P.G. MBUTHIA 43–47

Biochemical and haematological changes in rats administered an aqueous extract of *Prunus africana* stem bark at various dosage levels—P.K. GATHUMBI, J.W. MWANGI, S.M. NJIRO and G.M. MUGERA 123–128

GERBER, H.M.

Oral infectivity of cryopreserved *Haemonchus contortus* infective larvae that recovered relatively slowly when thawed after more than 15 years in liquid nitrogen—J.A. VAN WYK and H.M. GERBER 149–152

GERDES, G.H.

The prevalence of different African horsesickness virus serotypes in the Onderstepoort area near Pretoria, during an outbreak of African horsesickness in South Africa in 1995/1996—C.W. BREMER, G.H. GERDES, H. AITCHISON, I. LOUW, R.R. GREYLING and J. WELGEMOED 65–70

GREYLING, R.R.

The prevalence of different African horsesickness virus serotypes in the Onderstepoort area near Pretoria, during an outbreak of African horsesickness in South Africa in 1995/1996—C.W. BREMER, G.H. GERDES, H. AITCHISON, I. LOUW, R.R. GREYLING and J. WELGEMOED 65–70

GROBLER, D.G.

The rise and fall of tuberculosis in a free-ranging chacma baboon troop in the Kruger National Park—D.F. KEET, N.P.J. KRIEK, R.G. BENGIS, D.G. GROBLER and A. MICHEL 115–122

GROENEWALD, H.B.

Morphological changes associated with the development of the rumino-reticulum in growing lambs fed different rations—G.E. SWAN and H.B. GROENEWALD 105–114

HENTON, M.M.

Aerobic intestinal flora of wild-caught African dwarf crocodiles *Osteolaemus tetraspis*—F.W. HUCHZERMEYER, M.M. HENTON, J. RILEY and M. AGNAGNA 201–204

HERING-HAGENBECK, S.F.B.N.

A check-list of the nematode parasites of South African serpentes (snakes) and sauria (lizards)—S.F.B.N. HERING-HAGENBECK and J. BOOMKER 1–13

HIRUMI, H.

Suggested dosage rates of melarsoprol in the treatment of mice experimentally infected with *Trypanosoma brucei gambiense*—P.A. MBATI, K. HIRUMI, N. INOUE, N.H. SITUAKIBANZA and H. HIRUMI 71–74

HIRUMI, K.

Suggested dosage rates of melarsoprol in the treatment of mice experimentally infected with *Trypanosoma brucei gambiense*—P.A. MBATI, K. HIRUMI, N. INOUE, N.H. SITUAKIBANZA and H. HIRUMI 71–74

HORAK, I.G.

Parasites of South African wildlife. XVI. Helminths of some antelope species from the Eastern and Western Cape Provinces—J. BOOMKER, I.G. HORAK, R. WATERMEYER and D.G. BOOYSE 31–41

Parasites of domestic and wild animals in South Africa. XXXVIII. Ixodid ticks collected from 23 wild carnivore species—I.G. HORAK, L.E.O. BRAACK, L.J. FOURIE and JANE B. WALKER 239–250

HUCHZERMEYER, F.W.

Aerobic intestinal flora of wild-caught African dwarf crocodiles *Osteolaemus tetraspis*—F.W. HUCHZERMEYER, M.M. HENTON, J. RILEY and M. AGNAGNA 201–204

INOUE, N.

Suggested dosage rates of melarsoprol in the treatment of mice experimentally infected with *Trypanosoma brucei gambiense*—P.A. MBATI, K. HIRUMI, N. INOUE, N.H. SITUAKIBANZA and H. HIRUMI 71–74

JAMAL, S.

The distribution of tsetse (Diptera: Glossinidae) and bovine trypanosomosis in the Matutuine District, Maputo Province, Mozambique—I. SIGAUQUE, P. VAN DEN BOSSCHE, M. MOIANA, S. JAMAL and L. NEVES 167–172

JOSLING, D.

Ultrastructural study of the luminal surface of the ducts of the epididymis of gallinaceous birds—T.A. AIRE and D. JOSLING 191–199

KAGIRA, J.

The role of parasitic diseases as causes of mortality in cattle in a high potential area of central Kenya: a quantitative analysis—P.W.N. KANYARI and J. KAGIRA 157–161

KANYARI, P.W.N.

The role of parasitic diseases as causes of mortality in cattle in a high potential area of central Kenya: a quantitative analysis—P.W.N. KANYARI and J. KAGIRA 157–161

KAPPMEIER, KARIN

A newly developed odour-baited "H trap" for the live collection of *Glossina brevipalpis* and *Glossina austeni* (Diptera: Glossinidae) in South Africa—KARIN KAPPMEIER 15–26

Diurnal activity patterns of *Glossina brevipalpis* and *G. austeni* (Diptera: Glossinidae) in South Africa, with reference to season and meteorological factors—KARIN KAPPMEIER 179–189

KEET, D.F.

The rise and fall of tuberculosis in a free-ranging chacma baboon troop in the Kruger National Park—D.F. KEET, N.P.J. KRIEK, R.G. BENGIS, D.G. GROBLER and A. MICHEL 115–122

KHALIL, L.F.

Observations on a lesser-known monogenean, *Udonella myliobati*, from a copepod parasite, *Lepeophtheirus natalensis*, parasitizing the spotted ragged-tooth shark, *Carcharias taurus*, from South African waters—P.A.S. OLIVIER, S.M. DIPPENAAR, L.F. KHALIL and N.M. MOKGALONG 135–140

KOKOSHAROV, T.

Bacterial colonization and endotoxin activity during experimental acute fowl typhoid in chickens—T. KOKOSHAROV . 297–300

KRIEK, N.P.J.

The rise and fall of tuberculosis in a free-ranging chacma baboon troop in the Kruger National Park—D.F. KEET, N.P.J. KRIEK, R.G. BENGIS, D.G. GROBLER and A. MICHEL 115–122

Selected chemical parameters in the blood and metals in the organs of the Nile crocodile, *Crocodylus niloticus*, in the Kruger National Park—D. SWANEPOEL, J. BOOMKER and N.P.J. KRIEK 141–148

LEVINS, R.

Models for regional heartwater epidemiology in a variable environment—R. LEVINS 163–165

LOSHO, T.C.

Bovine borreliosis in Botswana—S.P. SHARMA, W. AMANFU and T.C. LOSHO 221–223

LOUW, I.

The prevalence of different African horsesickness virus serotypes in the Onderstepoort area near Pretoria, during an outbreak of African horsesickness in South Africa in 1995/1996—C.W. BREMER, G.H. GERDES, H. AITCHISON, I. LOUW, R.R. GREYLING and J. WELGEMOED 65–70

MABIKACHECHE, L.G.

Evaluation of a commercially available ELISA kit for detection of antibodies to *Anaplasma marginale* and *Anaplasma centrale* in cattle in Australia and Zimbabwe—P.M. BOWLES, J.B. MOLLOY, G.W. BLIGHT, S. SINGH and L.G. MABIKACHECHE 83–86

MAGAMBO, J.K.

Application of ultrasonography in prevalence studies of hydatid cysts in goats in north-western Turkana, Kenya and Toposaland, southern Sudan—E.M. NJOROGE, P.M.F. MBITHI, J.M. GATHUMA, T.M. WACHIRA, J.K. MAGAMBO and E. ZEYHLE 251–255

MALAN, F.S.

Molineus cati n. sp. (Nematoda, Trichostrongylina, Molineoidea), a parasite of feral cats, *Felis catus* Linnaeus, 1758 in South Africa—MARIE-CLAUDE DURETTE-DESSET, J. BOOMKER and F.S. MALAN 173–177

MARÉ, L.

Phage types of *Salmonella enterica* serovar Enteritidis isolated in South Africa from 1991–1995—L. MARÉ, M.L. VAN DER WALT and L.M.T. DICKS 129–133

MASHEGO, S.N.

Occurrence of *Neodiplozoon polycotyleus* Paperna, 1973 (Diplozoidae: Monogenea) in cyprinid fish in South Africa—S.N. MASHEGO 153–154

MBATI, P.A.

Leishmania major infections in *Phlebotomus duboscqi* fed on murine models immunized with *L. major* sub-cellular antigens and sandfly gut antigens—P.A. MBATI, C.O. ANJILI, S. ODONGO, P. OGAJA and W. TONU 57–63

Suggested dosage rates of melarsoprol in the treatment of mice experimentally infected with *Trypanosoma brucei gambiense*—P.A. MBATI, K. HIRUMI, N. INOUE, N.H. SITUAKIBANZA and H. HIRUMI 71–74

MBITHI, P.M.F.

Application of ultrasonography in prevalence studies of hydatid cysts in goats in north-western Turkana, Kenya and Toposaland, southern Sudan—E.M. NJOROGE, P.M.F. MBITHI, J.M. GATHUMA, T.M. WACHIRA, J.K. MAGAMBO and E. ZEYHLE 251–255

MBUTHIA, P.G.

Suspected sheep-associated malignant catarrhal fever in a zero-grazed dairy herd in Kenya—C.M MULEI, P.K. GATHUMBI and P.G. MBUTHIA 43–47

McCRINDLE, R.I.

Investigation of biological samples for monofluoroacetate and *Dichapetalum cymosum* poisoning in southern Africa—P.P. MINNAAR, R.I. McCRINDLE, T.W. NAUDÉ and C.J. BOTHA 27–30

MEKONNEN, S.

Efficacy of flumethrin 1 % pour-on against ticks on cattle under field condition in Ethiopia—S. MEKONNEN 235–237

MICHEL, A.

The rise and fall of tuberculosis in a free-ranging chacma baboon troop in the Kruger National Park—D.F. KEET, N.P.J. KRIEK, R.G. BENGIS, D.G. GROBLER and A. MICHEL 115–122

MINNAAR, P.P.

Investigation of biological samples for monofluoroacetate and *Dichapetalum cymosum* poisoning in southern Africa—P.P. MINNAAR, R.I. McCRINDLE, T.W. NAUDÉ and C.J. BOTHA 27–30

MOHAN, K.

Capsular serogroups of *Pasteurella multocida* isolated from animals in Zimbabwe—F. DZIVA, K. MOHAN and A. PAWANDIWA 225–228

Pasteurella gallinarum: Zimbabwean experience of a versatile pathogen—K. MOHAN, F. DZIVA and D. CHITAURO 301–305

MOIANA, M.

The distribution of tsetse (Diptera: Glossinidae) and bovine trypanosomosis in the Matutuine District, Maputo Province, Mozambique—I. SIGAUQUE, P. VAN DEN BOSSCHE, M. MOIANA, S. JAMAL and L. NEVES 167–172

MOKGALONG, N.M.

Observations on a lesser-known monogenean, *Udonella myliobati*, from a copepod parasite, *Lepeophtheirus natalensis*, parasitizing the spotted ragged-tooth shark, *Carcharias taurus*, from South African waters—P.A.S. OLIVIER, S.M. DIPPENAAR, L.F. KHALIL and N.M. MOKGALONG 135–140

MOLLOY, J.B.

Evaluation of a commercially available ELISA kit for detection of antibodies to *Anaplasma marginale* and *Anaplasma centrale* in cattle in Australia and Zimbabwe—P.M. BOWLES, J.B. MOLLOY, G.W. BLIGHT, S. SINGH and L.G. MABIKACHECHE 83–86

MOSALA, K.P.

The prevalence of internal and external parasites in pigs of different ages and sexes in Southeast District, Botswana—S.J. NSOSO, K.P. MOSALA, R.T. NDEBELE and S.S. RAMABU 217–220

MUGERA, G.M.

Biochemical and haematological changes in rats administered an aqueous extract of *Prunus africana* stem bark at various dosage levels—P.K. GATHUMBI, J.W. MWANGI, S.M. NJIRO and G.M. MUGERA 123–128

MÜLDERS, MARIA S.G.

Stability and partitioning of closantel and rafoxanide in ruminal fluid of sheep—G.E. SWAN, D.W. OLIVER, J. VAN RENSBURG, H.S. STEYN and MARIA S.G. MÜLDERS 97–103

MULEI, C.M.

Suspected sheep-associated malignant catarrhal fever in a zero-grazed dairy herd in Kenya—C.M MULEI, P.K. GATHUMBI and P.G. MBUTHIA 43–47

MUSHI, E.Z.

Helminth parasites of domestic pigeons (*Columba livia domestica*) in Sebele, Gaborone, Botswana—E.Z. MUSHI, M.G. BINTA, R.G. CHABO, R. NDEBELE and R. PANZIRAH 75–76

MWANGI, J.W.

Biochemical and haematological changes in rats administered an aqueous extract of *Prunus africana* stem bark at various dosage levels—P.K. GATHUMBI, J.W. MWANGI, S.M. NJIRO and G.M. MUGERA 123–128

NAUDÉ, T.W.

Investigation of biological samples for monofluoroacetate and *Dichapetalum cymosum* poisoning in southern Africa—P.P. MINNAAR, R.I. McCRINDLE, T.W. NAUDÉ and C.J. BOTHA 27–30

NDEBELE, R.

Helminth parasites of domestic pigeons (*Columba livia domestica*) in Sebele, Gaborone, Botswana—E.Z. MUSHI, M.G. BINTA, R.G. CHABO, R. NDEBELE and R. PANZIRAH 75–76

NDEBELE, R.T.

The prevalence of internal and external parasites in pigs of different ages and sexes in Southeast District, Botswana—S.J. NSOSO, K.P. MOSALA, R.T. NDEBELE and S.S. RAMABU 217–220

NEVES, L.

The distribution of tsetse (Diptera: Glossinidae) and bovine trypanosomosis in the Matutuine District, Maputo Province, Mozambique—I. SIGAUQUE, P. VAN DEN BOSSCHE, M. MOIANA, S. JAMAL and L. NEVES 167–172

NJIRO, S.M.

Biochemical and haematological changes in rats administered an aqueous extract of *Prunus africana* stem bark at various dosage levels—P.K. GATHUMBI, J.W. MWANGI, S.M. NJIRO and G.M. MUGERA 123–128

NJOROGE, E.M.

Application of ultrasonography in prevalence studies of hydatid cysts in goats in north-western Turkana, Kenya and Toposaland, southern Sudan—E.M. NJOROGE, P.M.F. MBITHI, J.M. GATHUMA, T.M. WACHIRA, J.K. MAGAMBO and E. ZEYHLE 251–255

NSOSO, S.J.

The prevalence of internal and external parasites in pigs of different ages and sexes in Southeast District, Botswana—S.J. NSOSO, K.P. MOSALA, R.T. NDEBELE and S.S. RAMABU 217–220

ODENDAAL, M.W.

The influence of *Mannheimia haemoloytica* A1 seed culture inoculum cell density on the production of leukotoxin in submerged culture supernatant—M.W. ODENDAAL and L. DU PLESSIS 205–216

ODONGO, S.

Leishmania major infections in *Phlebotomus duboscqi* fed on murine models immunized with *L. major* sub-cellular antigens and sandfly gut antigens—P.A. MBATI, C.O. ANJILI, S. ODONGO, P. OGAJA and W. TONU 57–63

OGAJA, P.

Leishmania major infections in *Phlebotomus duboscqi* fed on murine models immunized with *L. major* sub-cellular antigens and sandfly gut antigens—P.A. MBATI, C.O. ANJILI, S. ODONGO, P. OGAJA and W. TONU 57–63

OLIVER, D.W.

Stability and partitioning of closantel and rafoxanide in ruminal fluid of sheep—G.E. SWAN, D.W. OLIVER, J. VAN RENSBURG, H.S. STEYN and MARIA S.G. MÜLDERS 97–103

OLIVIER, P.A.S.

Observations on a lesser-known monogenean, *Udonella myliobati*, from a copepod parasite, *Lepeophtheirus natalensis*, parasitizing the spotted ragged-tooth shark, *Carcharias taurus*, from South African waters—P.A.S. OLIVIER, S.M. DIPPENAAR, L.F. KHALIL and N.M. MOKGALONG 135–140

OLUWANIYI, J.O.

Differential expression of surface membrane antigens on bovine monocytes activated with recombinant cytokines and during *Trypanosoma congolense* infection—V.O. TAIWO, V.O. ANOSA and J.O. OLUWANIYI 289–296

PANZIRAH, R.

Helminth parasites of domestic pigeons (*Columba livia domestica*) in Sebele, Gaborone, Botswana—E.Z. MUSHI, M.G. BINTA, R.G. CHABO, R. NDEBELE and R. PANZIRAH 75–76

PAWANDIWA, A.

Capsular serogroups of *Pasteurella multocida* isolated from animals in Zimbabwe—F. DZIVA, K. MOHAN and A PAWANDIWA, A. 225–228

PUTTERILL, J.F.

Oribatid mites (Acari, Oribatida) as intermediate hosts of tapeworms of the Family Anoplocephalidae (Cestoda) and the transmission of *Moniezia expansa* cysticercoids in South Africa—R. SCHUSTER, L. COETZEE and J.F. PUTTERILL 49–55

Studies on the genus *Setaria* Viborg, 1795 in South Africa. I. *Setaria africana* (Yeh, 1959)—R. WATERMEYER, J. BOOMKER and J.F. PUTTERILL 229–234

RAMABU, S.S.

The prevalence of internal and external parasites in pigs of different ages and sexes in Southeast District, Botswana—S.J. NSOSO, K.P. MOSALA, R.T. NDEBELE and S.S. RAMABU 217–220

RILEY, J.

Aerobic intestinal flora of wild-caught African dwarf crocodiles *Osteolaemus tetraspis*—F.W. HUCHZERMEYER, M.M. HENTON, J. RILEY and M. AGNAGNA 201–204

SCHUSTER, L.

Oribatid mites (Acari, Oribatida) as intermediate hosts of tapeworms of the Family Anoplocephalidae (Cestoda) and the transmission of *Moniezia expansa* cysticercoids in South Africa—R. SCHUSTER, L. COETZEE and J.F. PUTTERILL 49–55

SEWANI-RUSIKE, C.R.

Fertility in rats immunized with steroid-free bovine follicular fluid—C.R. SEWANI-RUSIKE and C. DAKWA 257–262

SHARMA, S.P.

Bovine borreliosis in Botswana—S.P. SHARMA, W. AMANFU and T.C. LOSHO 221–223

SIGAUQUE, I.

The distribution of tsetse (Diptera: Glossinidae) and bovine trypanosomosis in the Matutuine District, Maputo Province, Mozambique—I. SIGAUQUE, P. VAN DEN BOSSCHE, M. MOIANA, S. JAMAL and L. NEVES 167–172

SINGH, S.

Evaluation of a commercially available ELISA kit for detection of antibodies to *Anaplasma marginale* and *Anaplasma centrale* in cattle in Australia and Zimbabwe—P.M. BOWLES, J.B. MOLLOY, G.W. BLIGHT, S. SINGH and L.G. MABIKACHECHE 83–86

SITUAKIBANZA, N.H.

Suggested dosage rates of melarsoprol in the treatment of mice experimentally infected with *Trypanosoma brucei gambiense*—P.A. MBATI, K. HIRUMI, N. INOUE, N.H. SITUAKIBANZA and H. HIRUMI 71–74

STEYN, H.S.

Stability and partitioning of closantel and rafoxanide in ruminal fluid of sheep—G.E. SWAN, D.W. OLIVER, J. VAN RENSBURG, H.S. STEYN and MARIA S.G. MÜLDERS 97–103

SWAN, G.E.

Stability and partitioning of closantel and rafoxanide in ruminal fluid of sheep—G.E. SWAN, D.W. OLIVER, J. VAN RENSBURG, H.S. STEYN and MARIA S.G. MÜLDERS 97–103

Morphological changes associated with the development of the rumino-reticulum in growing lambs fed different rations—G.E. SWAN and H.B. GROENEWALD 105–114

SWANEPOEL, J.

Selected chemical parameters in the blood and metals in the organs of the Nile crocodile, *Crocodylus niloticus*, in the Kruger National Park—D. SWANEPOEL, J. BOOMKER and N.P.J. KRIEK 141–148

TAIWO, V.O.

In vitro erythrophagocytosis by cultured macrophages stimulated with extraneous substances and those isolated from the blood, spleen and bone marrow of Boran and N'Dama cattle infected with *Trypanosoma congolense* and *T. vivax*—V.O. TAIWO and V.O. ANOSA 273–287

Differential expression of surface membrane antigens on bovine monocytes activated with recombinant cytokines and during *Trypanosoma congolense* infection—V.O. TAIWO, V.O. ANOSA and J.O. OLUWANIYI 289–296

TONU, W.

Leishmania major infections in *Phlebotomus duboscqi* fed on murine models immunized with *L. major* sub-cellular antigens and sandfly gut antigens—P.A. MBATI, C.O. ANJILI, S. ODONGO, P. OGAJA and W. TONU 57–63

VAN DEN BOSSCHE, P.

The distribution of tsetse (Diptera: Glossinidae) and bovine trypanosomosis in the Matutuine District, Maputo Province, Mozambique—I. SIGAUQUE, P. VAN DEN BOSSCHE, M. MOIANA, S. JAMAL and L. NEVES 167–172

VAN DER WALT, M.L.

Phage types of *Salmonella enterica* serovar Enteritidis isolated in South Africa from 1991–1995—L. MARÉ, M.L. VAN DER WALT and L.M.T. DICKS 129–133

VAN RENSBURG, J.

Stability and partitioning of closantel and rafoxanide in ruminal fluid of sheep—G.E. SWAN, D.W. OLIVER, J. VAN RENSBURG, H.S. STEYN and MARIA S.G. MÜLDERS 97–103

VAN WYK, J.A.

Oral infectivity of cryopreserved *Haemonchus contortus* infective larvae that recovered relatively slowly when thawed after more than 15 years in liquid nitrogen—J.A. VAN WYK and H.M. GERBER 149–152

VERWOERD, D.W.

The effect of a natural maedi-visna virus infection on the productivity of South African sheep—B. DUNGU, J. VORSTER, G.F. BATH and D.W. VERWOERD 87–96

VORSTER, J.

The effect of a natural maedi-visna virus infection on the productivity of South African sheep—B. DUNGU, J. VORSTER, G.F. BATH and D.W. VERWOERD 87–96

WACHIRA, T.M.

Application of ultrasonography in prevalence studies of hydatid cysts in goats in north-western Turkana, Kenya and Toposaland, southern Sudan—E.M. NJOROGE, P.M.F. MBITHI, J.M. GATHUMA, T.M. WACHIRA, J.K. MAGAMBO and E. ZEYHLE 251–255

WALKER, JANE B.

Parasites of domestic and wild animals in South Africa. XXXVIII. Ixodid ticks collected from 23 wild carnivore species—I.G. HORAK, L.E.O. BRAACK, L.J. FOURIE and JANE B. WALKER 239–250

WATERMEYER, R.

Studies on the genus *Setaria* Viborg, 1795 in South Africa. I. *Setaria africana* (Yeh, 1959)—R. WATERMEYER, J. BOOMKER and J.F. PUTTERILL 229–234

Parasites of South African wildlife. XVI. Helminths of some antelope species from the Eastern and Western Cape Provinces—J. BOOMKER, I.G. HORAK, R. WATERMEYER and D.G. BOOYSE 31–41

WELGEMOED, J.

The prevalence of different African horsesickness virus serotypes in the Onderstepoort area near Pretoria, during an outbreak of African horsesickness in South Africa in 1995/1996—C.W. BREMER, G.H. GERDES, H. AITCHISON, I. LOUW, R.R. GREYLING and J. WELGEMOED 65–70

ZEYHLE, E.

Application of ultrasonography in prevalence studies of hydatid cysts in goats in north-western Turkana, Kenya and Toposaland, southern Sudan—E.M. NJOROGE, P.M.F. MBITHI, J.M. GATHUMA, T.M. WACHIRA, J.K. MAGAMBO and E. ZEYHLE 251–255

ONDERSTEPOORT JOURNAL OF VETERINARY RESEARCH

Volume 67 • Numbers 1–4 • 2000

Subject index

Abbreviata

A check-list of the nematode parasites of South African serpentes (snakes) and sauria (lizards)—S.F.B.N. HERINGHAGENBECK and J. BOOMKER 1–13

Acari

Oribatid mites (Acari, Oribatida) as intermediate hosts of tapeworms of the Family Anoplocephalidae (Cestoda) and the transmission of *Moniezia expansa* cysticercoids in South Africa—R. SCHUSTER, L. COETZEE and J.F. PUTTERILL 49–55

Acaricide

Efficacy of flumethrin 1 % pour-on against ticks on cattle under field condition in Ethiopia—S. MEKONNEN 235–237

Aeromonas hydrophila

Aerobic intestinal flora of wild-caught African dwarf crocodiles *Osteolaemus tetraspis*—F.W. HUCHZERMEYER, M.M. HENTON, J. RILEY and M. AGNAGNA 201–204

African horsesickness virus

The prevalence of different African horsesickness virus serotypes in the Onderstepoort area near Pretoria, during an outbreak of African horsesickness in South Africa in 1995/1996—C.W. BREMER, G.H. GERDES, H. AITCHISON, I. LOUW, R.R. GREYLING and J. WELGEMOED 65–70

African dwarf crocodile

Aerobic intestinal flora of wild-caught African dwarf crocodiles *Osteolaemus tetraspis*—F.W. HUCHZERMEYER, M.M. HENTON, J. RILEY and M. AGNAGNA 201–204

Agglutination

Evaluation of a commercially available ELISA kit for detection of antibodies to *Anaplasma marginale* and *Anaplasma centrale* in cattle in Australia and Zimbabwe—P.M. BOWLES, J.B. MOLLOY, G.W. BLIGHT, S. SINGH and L.G. MABIKACHECHE 83–86

Alcelaphus buselaphus

Parasites of South African wildlife. XVI. Helminths of some antelope species from the Eastern and Western Cape Provinces—J. BOOMKER, I.G. HORAK, R. WATERMEYER and D.G. BOOYSE 31–41

Anaemia

In vitro erythrophagocytosis by cultured macrophages stimulated with extraneous substances and those isolated from the blood, spleen and bone marrow of Boran and N'Dama cattle infected with *Trypanosoma congolense* and *Trypanosoma vivax*—V.O. TAIWO and V.O. ANOSA 273–287

Differential expression of surface membrane antigens on bovine monocytes activated with recombinant cytokines and during *Trypanosoma congolense* infection—V.O. TAIWO, V.O. ANOSA and J.O. OLUWANIYI 289–296

Anaplasma centrale, Anaplasma marginale

Evaluation of a commercially available ELISA kit for detection of antibodies to *Anaplasma marginale* and *Anaplasma centrale* in cattle in Australia and Zimbabwe—P.M. BOWLES, J.B. MOLLOY, G.W. BLIGHT, S. SINGH and L.G. MABIKACHECHE 83–86

Anaplasmosis
The role of parasitic diseases as causes of mortality in cattle in a high potential area of central Kenya: a quantitative analysis—P.W.N. KANYARI and J. KAGIRA 157–161

Anoplocephalidae
Oribatid mites (Acari, Oribatida) as intermediate hosts of tapeworms of the Family Anoplocephalidae (Cestoda) and the transmission of *Moniezia expansa* cysticercoids in South Africa—R. SCHUSTER, L. COETZEE and J.F. PUTTERILL 49–55

Antelope parasites
Parasites of South African wildlife. XVI. Helminths of some antelope species from the Eastern and Western Cape Provinces—J. BOOMKER, I.G. HORAK, R. WATERMEYER and D.G. BOOYSE 31–41

Antibodies
Evaluation of a commercially available ELISA kit for detection of antibodies to *Anaplasma marginale* and *Anaplasma centrale* in cattle in Australia and Zimbabwe—P.M. BOWLES, J.B. MOLLOY, G.W. BLIGHT, S. SINGH and L.G. MABIKACHECHE 83–86

Antidorcas marsupialis
Parasites of South African wildlife. XVI. Helminths of some antelope species from the Eastern and Western Cape Provinces—J. BOOMKER, I.G. HORAK, R. WATERMEYER and D.G. BOOYSE 31–41

Antigen
Leishmania major infections in *Phlebotomus duboscqi* fed on murine models immunized with *L. major* sub-cellular antigens and sandfly gut antigens—P.A. MBATI, C.O. ANJILI, S. ODONGO, P. OGAJA and W. TONU 57–63

Differential expression of surface membrane antigens on bovine monocytes activated with recombinant cytokines and during *Trypanosoma congolense* infection—V.O. TAIWO, V.O. ANOSA and J.O. OLUWANIYI 289–296

Ascardia columbae
Helminth parasites of domestic pigeons (*Columba livia domestica*) in Sebele, Gaborone, Botswana—E.Z. MUSHI, M.G. BINTA, R.G. CHABO, R. NDEBELE and R. PANZIRAH 75–76

Ascaris suum
The prevalence of internal and external parasites in pigs of different ages and sexes in Southeast District, Botswana—S.J. NSOSO, K.P. MOSALA, R.T. NDEBELE and S.S. RAMABU 217–220

Babesia bovis
Bovine borreliosis in Botswana—S.P. SHARMA, W. AMANFU and T.C. LOSHO 221–223

Babesiosis
The role of parasitic diseases as causes of mortality in cattle in a high potential area of central Kenya: a quantitative analysis—P.W.N. KANYARI and J. KAGIRA 157–161

Baboon tuberculosis
The rise and fall of tuberculosis in a free-ranging chacma baboon troop in the Kruger National Park—D.F. KEET, N.P.J. KRIEK, R.G. BENGIS, D.G. GROBLER and A. MICHEL 115–122

Bacterial colonization
Bacterial colonization and endotoxin activity during experimental acute fowl typhoid in chickens—T. KOKOSHAROV . 297–300

***Barbus* spp.**
Occurrence of *Neodiplozoon polycotyleus* Paperna, 1973 (Diplozoidae: Monogenea) in cyprinid fish in South Africa—S.N. MASHEGO 153–154

Bayticol pour-on
Efficacy of flumethrin 1 % pour-on against ticks on cattle under field condition in Ethiopia—S. MEKONNEN 235–237

Birds
Ultrastructural study of the luminal surface of the ducts of the epididymis of gallinaceous birds—T.A. AIRE and D. JOSLING 191–199

Bitis arietans
A check-list of the nematode parasites of South African serpentes (snakes) and sauria (lizards)—S.F.B.N. HERINGHAGENBECK and J. BOOMKER 1–13

Blood
Selected chemical parameters in the blood and metals in the organs of the Nile crocodile, *Crocodylus niloticus*, in the Kruger National Park—D. SWANEPOEL, J. BOOMKER and N.P.J. KRIEK 141–148

In vitro erythrophagocytosis by cultured macrophages stimulated with extraneous substances and those isolated from the blood, spleen and bone marrow of Boran and N'Dama cattle infected with *Trypanosoma congolense* and *Trypanosoma vivax*—V.O. TAIWO and V.O. ANOSA 273–287

Bone marrow

In vitro erythrophagocytosis by cultured macrophages stimulated with extraneous substances and those isolated from the blood, spleen and bone marrow of Boran and N'Dama cattle infected with *Trypanosoma congolense* and *Trypanosoma vivax*—V.O. TAIWO and V.O. ANOSA 273–287

Boran cattle

In vitro erythrophagocytosis by cultured macrophages stimulated with extraneous substances and those isolated from the blood, spleen and bone marrow of Boran and N'Dama cattle infected with *Trypanosoma congolense* and *Trypanosoma vivax*—V.O. TAIWO and V.O. ANOSA 273–287

Differential expression of surface membrane antigens on bovine monocytes activated with recombinant cytokines and during *Trypanosoma congolense* infection—V.O. TAIWO, V.O. ANOSA and J.O. OLUWANIYI 289–296

Borrelia theileri

Bovine borreliosis in Botswana—S.P. SHARMA, W. AMANFU and T.C. LOSHO 221–223

Botswana

Helminth parasites of domestic pigeons (*Columba livia domestica*) in Sebele, Gaborone, Botswana—E.Z. MUSHI, M.G. BINTA, R.G. CHABO, R. NDEBELE and R. PANZIRAH 75–76

The prevalence of internal and external parasites in pigs of different ages and sexes in Southeast District, Botswana—S.J. NSOSO, K.P. MOSALA, R.T. NDEBELE and S.S. RAMABU 217–220

Bovine borreliosis in Botswana—S.P. SHARMA, W. AMANFU and T.C. LOSHO 221–223

Bovine

The distribution of tsetse (Diptera: Glossinidae) and bovine trypanosomosis in the Matutuine District, Maputo Province, Mozambique—I. SIGAUQUE, P. VAN DEN BOSSCHE, M. MOIANA, S. JAMAL and L. NEVES 167–172

Bovine borreliosis in Botswana—S.P. SHARMA, W. AMANFU and T.C. LOSHO 221–223

Differential expression of surface membrane antigens on bovine monocytes activated with recombinant cytokines and during *Trypanosoma congolense* infection—V.O. TAIWO, V.O. ANOSA and J.O. OLUWANIYI 289–296

Capsular serogroups

Capsular serogroups of *Pasteurella multocida* isolated from animals in Zimbabwe—F. DZIVA, K. MOHAN and A. PAWANDIWA 225–228

Carcharias taurus

Observations on a lesser-known monogenean, *Udonella myliobati*, from a copepod parasite, *Lepeophtheirus natalensis*, parasitizing the spotted ragged-tooth shark, *Carcharias taurus*, from South African waters—P.A.S. OLIVIER, S.M. DIPPENAAR, L.F. KHALIL and N.M. MOKGALONG 135–140

Carnivore ticks

Parasites of domestic and wild animals in South Africa. XXXVIII. Ixodid ticks collected from 23 wild carnivore species—I.G. HORAK, L.E.O. BRAACK, L.J. FOURIE and JANE B. WALKER 239–250

Catarrhal fever

Suspected sheep-associated malignant catarrhal fever in a zero-grazed dairy herd in Kenya—C.M MULEI, P.K. GATHUMBI and P.G. MBUTHIA 43–47

Cats

Molineus cati n. sp. (Nematoda, Trichostrongylina, Molineoidea), a parasite of feral cats, *Felis catus* Linnaeus, 1758 in South Africa—MARIE-CLAUDE DURETTE-DESSET, J. BOOMKER and F.S. MALAN 173–177

Parasites of domestic and wild animals in South Africa. XXXVIII. Ixodid ticks collected from 23 wild carnivore species—I.G. HORAK, L.E.O. BRAACK, L.J. FOURIE and JANE B. WALKER 239–250

Cattle

Evaluation of a commercially available ELISA kit for detection of antibodies to *Anaplasma marginale* and *Anaplasma centrale* in cattle in Australia and Zimbabwe—P.M. BOWLES, J.B. MOLLOY, G.W. BLIGHT, S. SINGH and L.G. MABIKACHECHE 83–86

The role of parasitic diseases as causes of mortality in cattle in a high potential area of central Kenya: a quantitative analysis—P.W.N. KANYARI and J. KAGIRA 157–161

Efficacy of flumethrin 1 % pour-on against ticks on cattle under field condition in Ethiopia—S. MEKONNEN 235–237

Parameters for the determination and evaluation of heat stress in dairy cattle in South Africa—J.H. DU PREEZ 263–271

In vitro erythrophagocytosis by cultured macrophages stimulated with extraneous substances and those isolated from the blood, spleen and bone marrow of Boran and N'Dama cattle infected with *Trypanosoma congolense* and *Trypanosoma vivax*—V.O. TAIWO and V.O. ANOSA 273–287

Differential expression of surface membrane antigens on bovine monocytes activated with recombinant cytokines and during *Trypanosoma congolense* infection—V.O. TAIWO, V.O. ANOSA and J.O. OLUWANIYI 289–296

Cestoda

Oribatid mites (Acari, Oribatida) as intermediate hosts of tapeworms of the Family Anoplocephalidae (Cestoda) and the transmission of *Moniezia expansa* cysticercoids in South Africa—R. SCHUSTER, L. COETZEE and J.F. PUTTERILL 49–55

Chacma

The rise and fall of tuberculosis in a free-ranging chacma baboon troop in the Kruger National Park—D.F. KEET, N.P.J. KRIEK, R.G. BENGIS, D.G. GROBLER and A. MICHEL 115–122

Chamaeleo dilepis

A check-list of the nematode parasites of South African serpentes (snakes) and sauria (lizards)—S.F.B.N. HERINGHAGENBECK and J. BOOMKER 1–13

Chickens

Bacterial colonization and endotoxin activity during experimental acute fowl typhoid in chickens—T. KOKOSHAROV . 297–300

Pasteurella gallinarum: Zimbabwean experience of a versatile pathogen—K. MOHAN, F. DZIVA and D. CHITAURO 301–305

Chlamydia

Pasteurella gallinarum: Zimbabwean experience of a versatile pathogen—K. MOHAN, F. DZIVA and D. CHITAURO 301–305

Closantel

Stability and partitioning of closantel and rafoxanide in ruminal fluid of sheep—G.E. SWAN, D.W. OLIVER, J. VAN RENSBURG, H.S. STEYN and MARIA S.G. MÜLDERS 97–103

Columba livia domestica

Helminth parasites of domestic pigeons (*Columba livia domestica*) in Sebele, Gaborone, Botswana—E.Z. MUSHI, M.G. BINTA, R.G. CHABO, R. NDEBELE and R. PANZIRAH 75–76

Copepod

Observations on a lesser-known monogenean, *Udonella myliobati*, from a copepod parasite, *Lepeophtheirus natalensis*, parasitizing the spotted ragged-tooth shark, *Carcharias taurus*, from South African waters—P.A.S. OLIVIER, S.M. DIPPENAAR, L.F. KHALIL and N.M. MOKGALONG 135–140

Cortisol

Parameters for the determination and evaluation of heat stress in dairy cattle in South Africa—J.H. DU PREEZ 263–271

Coturnix coturnix

Ultrastructural study of the luminal surface of the ducts of the epididymis of gallinaceous birds—T.A. AIRE and D. JOSLING 191–199

Connochaetes gnou

Parasites of South African wildlife. XVI. Helminths of some antelope species from the Eastern and Western Cape Provinces—J. BOOMKER, I.G. HORAK, R. WATERMEYER and D.G. BOOYSE 31–41

Crocodile

Selected chemical parameters in the blood and metals in the organs of the Nile crocodile, *Crocodylus niloticus*, in the Kruger National Park—D. SWANEPOEL, J. BOOMKER and N.P.J. KRIEK 141–148

Aerobic intestinal flora of wild-caught African dwarf crocodiles *Osteolaemus tetraspis*—F.W. HUCHZERMEYER, M.M. HENTON, J. RILEY and M. AGNAGNA 201–204

Crocodylis niloticus

Selected chemical parameters in the blood and metals in the organs of the Nile crocodile, *Crocodylus niloticus*, in the Kruger National Park—D. SWANEPOEL, J. BOOMKER and N.P.J. KRIEK 141–148

Culture inoculum, supernatant

The influence of *Mannheimia haemoloytica* A1 seed culture inoculum cell density on the production of leukotoxin in sub-merged culture supernatant—M.W. ODENDAAL and L. DU PLESSIS 205–216

Cultured macrophages

In vitro erythrophagocytosis by cultured macrophages stimulated with extraneous substances and those isolated from the blood, spleen and bone marrow of Boran and N'Dama cattle infected with *Trypanosoma congolense* and *Trypanosoma vivax*—V.O. TAIWO and V.O. ANOSA 273–287

Cysticercoids

Oribatid mites (Acari, Oribatida) as intermediate hosts of tapeworms of the Family Anoplocephalidae (Cestoda) and the transmission of *Moniezia expansa* cysticercoids in South Africa—R. SCHUSTER, L. COETZEE and J.F. PUTTERILL 49–55

Cytokines

In vitro erythrophagocytosis by cultured macrophages stimulated with extraneous substances and those isolated from the blood, spleen and bone marrow of Boran and N'Dama cattle infected with *Trypanosoma congolense* and *Trypanosoma vivax*—V.O. TAIWO and V.O. ANOSA 273–287

Differential expression of surface membrane antigens on bovine monocytes activated with recombinant cytokines and during *Trypanosoma congolense* infection—V.O. TAIWO, V.O. ANOSA and J.O. OLUWANIYI 289–296

Cytotoxin

The influence of *Mannheimia haemoloytica* A1 seed culture inoculum cell density on the production of leukotoxin in submerged culture supernatant—M.W. ODENDAAL and L. DU PLESSIS 205–216

Dairy cattle

Suspected sheep-associated malignant catarrhal fever in a zero-grazed dairy herd in Kenya—C.M MULEI, P.K. GATHUMBI and P.G. MBUTHIA 43–47

Efficacy of flumethrin 1 % pour-on against ticks on cattle under field condition in Ethiopia—S. MEKONNEN 235–237

Parameters for the determination and evaluation of heat stress in dairy cattle in South Africa—J.H. DU PREEZ 263–271

Damaliscus pygargus dorcas

Parasites of South African wildlife. XVI. Helminths of some antelope species from the Eastern and Western Cape Provinces—J. BOOMKER, I.G. HORAK, R. WATERMEYER and D.G. BOOYSE 31–41

Dichapetalum cymosum

Investigation of biological samples for monofluoroacetate and *Dichapetalum cymosum* poisoning in southern Africa—P.P. MINNAAR, R.I. McCRINDLE, T.W. NAUDÉ and C.J. BOTHA 27–30

Diplozoidae

Occurrence of *Neodiplozoon polycotyleus* Paperna, 1973 (Diplozoidae: Monogenea) in cyprinid fish in South Africa—S.N. MASHEGO 153–154

Diptera

A newly developed odour-baited "H trap" for the live collection of *Glossina brevipalpis* and *Glossina austeni* (Diptera: Glossinidae) in South Africa—KARIN KAPPMEIER 15–26

The distribution of tsetse (Diptera: Glossinidae) and bovine trypanosomosis in the Matutuine District, Maputo Province, Mozambique—I. SIGAUQUE, P. VAN DEN BOSSCHE, M. MOIANA, S. JAMAL and L. NEVES 167–172

Diurnal activity patterns of *Glossina brevipalpis* and *G. austeni* (Diptera: Glossinidae) in South Africa, with reference to season and meteorological factors—KARIN KAPPMEIER 179–189

Dorper sheep

The effect of a natural maedi-visna virus infection on the productivity of South African sheep—B. DUNGU, J. VORSTER, G.F. BATH and D.W. VERWOERD 87–96

Drug stability

Stability and partitioning of closantel and rafoxanide in ruminal fluid of sheep—G.E. SWAN, D.W. OLIVER, J. VAN RENSBURG, H.S. STEYN and MARIA S.G. MÜLDERS 97–103

Ducts

Ultrastructural study of the luminal surface of the ducts of the epididymis of gallinaceous birds—T.A. AIRE and D. JOSLING 191–199

East Coast fever

The role of parasitic diseases as causes of mortality in cattle in a high potential area of central Kenya: a quantitative analysis—P.W.N. KANYARI and J. KAGIRA 157–161

Eggs

Phage types of *Salmonella enterica* serovar Enteritidis isolated in South Africa from 1991–1995—L. MARÉ, M.L. VAN DER WALT and L.M.T. DICKS 129–133

Observations on a lesser-known monogenean, *Udonella myliobati*, from a copepod parasite, *Lepeophtheirus natalensis*, parasitizing the spotted ragged-tooth shark, *Carcharias taurus*, from South African waters—P.A.S. OLIVIER, S.M. DIPPENAAR, L.F. KHALIL and N.M. MOKGALONG 135–140

ELISA

Evaluation of a commercially available ELISA kit for detection of antibodies to *Anaplasma marginale* and *Anaplasma centrale* in cattle in Australia and Zimbabwe—P.M. BOWLES, J.B. MOLLOY, G.W. BLIGHT, S. SINGH and L.G. MABIKACHECHE 83–86

Endotoxin

Bacterial colonization and endotoxin activity during experimental acute fowl typhoid in chickens—T. KOKOSHAROV . 297–300

Enteritidis

Phage types of *Salmonella enterica* serovar Enteritidis isolated in South Africa from 1991–1995—L. MARÉ, M.L. VAN DER WALT and L.M.T. DICKS 129–133

Epidemiology

Models for regional heartwater epidemiology in a variable environment—R. LEVINS 163–165

Epididymis

Ultrastructural study of the luminal surface of the ducts of the epididymis of gallinaceous birds—T.A. AIRE and D. JOSLING 191–199

Escherichia coli

Aerobic intestinal flora of wild-caught African dwarf crocodiles *Osteolaemus tetraspis*—F.W. HUCHZERMEYER, M.M. HENTON, J. RILEY and M. AGNAGNA 201–204

Ethiopia

Efficacy of flumethrin 1 % pour-on against ticks on cattle under field conditions in Ethiopia—S. MEKONNEN 235–237

Felis catus

Molineus cati n. sp. (Nematoda, Trichostrongylina, Molineoidea), a parasite of feral cats, *Felis catus* Linnaeus, 1758 in South Africa—MARIE-CLAUDE DURETTE-DESSET, J. BOOMKER and F.S. MALAN 173–177

Fertility

Fertility in rats immunized with steroid-free bovine follicular fluid—C.R. SEWANI-RUSIKE and C. DAKWA 257–262

Fever

Suspected sheep-associated malignant catarrhal fever in a zero-grazed dairy herd in Kenya—C.M MULEI, P.K. GATHUMBI and P.G. MBUTHIA 43–47

Fish

Occurrence of *Neodiplozoon polycotyleus* Paperna, 1973 (Diplozoidae: Monogenea) in cyprinid fish in South Africa—S.N. MASHEGO 153–154

Flagella antigen

Leishmania major infections in *Phlebotomus duboscqi* fed on murine models immunized with *L. major* sub-cellular antigens and sandfly gut antigens—P.A. MBATI, C.O. ANJILI, S. ODONGO, P. OGAJA and W. TONU 57–63

Flies

A newly developed odour-baited "H trap" for the live collection of *Glossina brevipalpis* and *Glossina austeni* (Diptera: Glossinidae) in South Africa—KARIN KAPPMEIER 15–26

Diurnal activity patterns of *Glossina brevipalpis* and *G. austeni* (Diptera: Glossinidae) in South Africa, with reference to season and meteorological factors—KARIN KAPPMEIER 179–189

Flora

Aerobic intestinal flora of wild-caught African dwarf crocodiles *Osteolaemus tetraspis*—F.W. HUCHZERMEYER, M.M. HENTON, J. RILEY and M. AGNAGNA 201–204

Flumethrin

Efficacy of flumethrin 1 % pour-on against ticks on cattle under field condition in Ethiopia—S. MEKONNEN 235–237

Fowl typhoid

Bacterial colonization and endotoxin activity during experimental acute fowl typhoid in chickens—T. KOKOSHAROV . 297–300

FSH

Fertility in rats immunized with steroid-free bovine follicular fluid—C.R. SEWANI-RUSIKE and C. DAKWA 257–262

Gallus gallus

Ultrastructural study of the luminal surface of the ducts of the epididymis of gallinaceous birds—T.A. AIRE and D. JOSLING 191–199

Gifblaar

Investigation of biological samples for monofluoroacetate and *Dichapetalum cymosum* poisoning in southern Africa—P.P. MINNAAR, R.I. McCRINDLE, T.W. NAUDÉ and C.J. BOTHA 27–30

Glossinidae, *Glossina austeni, Glossina brevipalpis*

A newly developed odour-baited "H trap" for the live collection of *Glossina brevipalpis* and *Glossina austeni* (Diptera: Glossinidae) in South Africa—KARIN KAPPMEIER 15–26

The distribution of tsetse (Diptera: Glossinidae) and bovine trypanosomosis in the Matutuine District, Maputo Province, Mozambique—I. SIGAUQUE, P. VAN DEN BOSSCHE, M. MOIANA, S. JAMAL and L. NEVES 167–172

Diurnal activity patterns of *Glossina brevipalpis* and *G. austeni* (Diptera: Glossinidae) in South Africa, with reference to season and meteorological factors—KARIN KAPPMEIER 179–189

Goats

Phage types of *Salmonella enterica* serovar Enteritidis isolated in South Africa from 1991–1995—L. MARÉ, M.L. VAN DER WALT and L.M.T. DICKS 129–133

Application of ultrasonography in prevalence studies of hydatid cysts in goats in north-western Turkana, Kenya and Toposaland, southern Sudan—E.M. NJOROGE, P.M.F. MBITHI, J.M. GATHUMA, T.M. WACHIRA, J.K. MAGAMBO and E. ZEYHLE 251–255

Guinea fowls

Pasteurella gallinarum: Zimbabwean experience of a versatile pathogen—K. MOHAN, F. DZIVA and D. CHITAURO 301–305

Haemonchus contortus

Oral infectivity of cryopreserved *Haemonchus contortus* infective larvae that recovered relatively slowly when thawed after more than 15 years in liquid nitrogen—J.A. VAN WYK and H.M. GERBER 149–152

Haemophilus

Pasteurella gallinarum: Zimbabwean experience of a versatile pathogen—K. MOHAN, F. DZIVA and D. CHITAURO 301–305

Heartwater

The role of parasitic diseases as causes of mortality in cattle in a high potential area of central Kenya: a quantitative analysis—P.W.N. KANYARI and J. KAGIRA 157–161

Models for regional heartwater epidemiology in a variable environment—R. LEVINS 163–165

Heat stress

Parameters for the determination and evaluation of heat stress in dairy cattle in South Africa—J.H. DU PREEZ 263–271

Helminths

Parasites of South African wildlife. XVI. Helminths of some antelope species from the Eastern and Western Cape Provinces—J. BOOMKER, I.G. HORAK, R. WATERMEYER and D.G. BOOYSE 31–41

Helminth parasites of domestic pigeons (*Columba livia domestica*) in Sebele, Gaborone, Botswana—E.Z. MUSHI, M.G. BINTA, R.G. CHABO, R. NDEBELE and R. PANZIRAH 75–76

Occurrence of *Neodiplozoon polycotyleus* Paperna, 1973 (Diplozoidae: Monogenea) in cyprinid fish in South Africa—S.N. MASHEGO 153–154

Studies on the genus *Setaria* Viborg, 1795 in South Africa. I. *Setaria africana* (Yeh, 1959)—R. WATERMEYER, J. BOOMKER and J.F. PUTTERILL 229–234

Horsesickness virus

The prevalence of different African horsesickness virus serotypes in the Onderstepoort area near Pretoria, during an outbreak of African horsesickness in South Africa in 1995/1996—C.W. BREMER, G.H. GERDES, H. AITCHISON, I. LOUW, R.R. GREYLING and J. WELGEMOED 65–70

H trap

A newly developed odour-baited "H trap" for the live collection of *Glossina brevipalpis* and *Glossina austeni* (Diptera: Glossinidae) in South Africa—KARIN KAPPMEIER 15–26

Hydatid cysts

Application of ultrasonography in prevalence studies of hydatid cysts in goats in north-western Turkana, Kenya and Toposaland, southern Sudan—E.M. NJOROGE, P.M.F. MBITHI, J.M. GATHUMA, T.M. WACHIRA, J.K. MAGAMBO and E. ZEYHLE 251–255

Hydatidosis

The role of parasitic diseases as causes of mortality in cattle in a high potential area of central Kenya: a quantitative analysis—P.W.N. KANYARI and J. KAGIRA 157–161

Immunization

Leishmania major infections in *Phlebotomus duboscqi* fed on murine models immunized with *L. major* sub-cellular antigens and sandfly gut antigens—P.A. MBATI, C.O. ANJILI, S. ODONGO, P. OGAJA and W. TONU 57–63

Immunoneutralization

Fertility in rats immunized with steroid-free bovine follicular fluid—C.R. SEWANI-RUSIKE and C. DAKWA 257–262

***In vitro* erythrophagocytosis**

In vitro erythrophagocytosis by cultured macrophages stimulated with extraneous substances and those isolated from the blood, spleen and bone marrow of Boran and N'Dama cattle infected with *Trypanosoma congolense* and *Trypanosoma vivax*—V.O. TAIWO and V.O. ANOSA 273–287

Infection

Leishmania major infections in *Phlebotomus duboscqi* fed on murine models immunized with *L. major* sub-cellular antigens and sandfly gut antigens—P.A. MBATI, C.O. ANJILI, S. ODONGO, P. OGAJA and W. TONU 57–63

The effect of a natural maedi-visna virus infection on the productivity of South African sheep—B. DUNGU, J. VORSTER, G.F. BATH and D.W. VERWOERD 87–96

Oral infectivity of cryopreserved *Haemonchus contortus* infective larvae that recovered relatively slowly when thawed after more than 15 years in liquid nitrogen—J.A. VAN WYK and H.M. GERBER 149–152

Inhibin

Fertility in rats immunized with steroid-free bovine follicular fluid—C.R. SEWANI-RUSIKE and C. DAKWA 257–262

Kalicephalus

A check-list of the nematode parasites of South African serpentes (snakes) and sauria (lizards)—S.F.B.N. HERING-HAGENBECK and J. BOOMKER 1–13

Kenya

Suspected sheep-associated malignant catarrhal fever in a zero-grazed dairy herd in Kenya—C.M MULEI, P.K. GATHUMBI and P.G. MBUTHIA 43–47

The role of parasitic diseases as causes of mortality in cattle in a high potential area of central Kenya: a quantitative analysis—P.W.N. KANYARI and J. KAGIRA 157–161

Application of ultrasonography in prevalence studies of hydatid cysts in goats in north-western Turkana, Kenya and Toposaland, southern Sudan—E.M. NJOROGE, P.M.F. MBITHI, J.M. GATHUMA, T.M. WACHIRA, J.K. MAGAMBO and E. ZEYHLE 251–255

Kruger National Park

The rise and fall of tuberculosis in a free-ranging chacma baboon troop in the Kruger National Park—D.F. KEET, N.P.J. KRIEK, R.G. BENGIS, D.G. GROBLER and A. MICHEL 115–122

Selected chemical parameters in the blood and metals in the organs of the Nile crocodile, *Crocodylus niloticus*, in the Kruger National Park—D. SWANEPOEL, J. BOOMKER and N.P.J. KRIEK 141–148

Lambs

The effect of a natural maedi-visna virus infection on the productivity of South African sheep—B. DUNGU, J. VORSTER, G.F. BATH and D.W. VERWOERD 87–96

Morphological changes associated with the development of the rumino-reticulum in growing lambs fed different rations—G.E. SWAN and H.B. GROENEWALD 105–114

Larvae

Oral infectivity of cryopreserved *Haemonchus contortus* infective larvae that recovered relatively slowly when thawed after more than 15 years in liquid nitrogen—J.A. VAN WYK and H.M. GERBER 149–152

Leishmania major

Leishmania major infections in *Phlebotomus duboscqi* fed on murine models immunized with *L. major* sub-cellular antigens and sandfly gut antigens—P.A. MBATI, C.O. ANJILI, S. ODONGO, P. OGAJA and W. TONU 57–63

Lepeophtheirus natalensis

Observations on a lesser-known monogenean, *Udonella myliobati*, from a copepod parasite, *Lepeophtheirus natalensis*, parasitizing the spotted ragged-tooth shark, *Carcharias taurus*, from South African waters—P.A.S. OLIVIER, S.M. DIPPENAAR, L.F. KHALIL and N.M. MOKGALONG 135–140

Leukotoxin

The influence of *Mannheimia haemoloytica* A1 seed culture inoculum cell density on the production of leukotoxin in submerged culture supernatant—M.W. ODENDAAL and L. DU PLESSIS 205–216

Liquid chromatography

Investigation of biological samples for monofluoroacetate and *Dichapetalum cymosum* poisoning in southern Africa—P.P. MINNAAR, R.I. McCRINDLE, T.W. NAUDÉ and C.J. BOTHA 27–30

Lizards

A check-list of the nematode parasites of South African serpentes (snakes) and sauria (lizards)—S.F.B.N. HERING-HAGENBECK and J. BOOMKER 1–13

Longevity

Oral infectivity of cryopreserved *Haemonchus contortus* infective larvae that recovered relatively slowly when thawed after more than 15 years in liquid nitrogen—J.A. VAN WYK and H.M. GERBER 149–152

Maedi-visna virus

The effect of a natural maedi-visna virus infection on the productivity of South African sheep—B. DUNGU, J. VORSTER, G.F. BATH and D.W. VERWOERD 87–96

Mannheimia haemolytica

The influence of *Mannheimia haemoloytica* A1 seed culture inoculum cell density on the production of leukotoxin in submerged culture supernatant—M.W. ODENDAAL and L. DU PLESSIS 205–216

Mel-B

Suggested dosage rates of melarsoprol in the treatment of mice experimentally infected with *Trypanosoma brucei gambiense*—P.A. MBATI, K. HIRUMI, N. INOUE, N.H. SITUAKIBANZA and H. HIRUMI 71–74

Melarsopral

Suggested dosage rates of melarsoprol in the treatment of mice experimentally infected with *Trypanosoma brucei gambiense*—P.A. MBATI, K. HIRUMI, N. INOUE, N.H. SITUAKIBANZA and H. HIRUMI 71–74

Metals

Selected chemical parameters in the blood and metals in the organs of the Nile crocodile, *Crocodylus niloticus*, in the Kruger National Park—D. SWANEPOEL, J. BOOMKER and N.P.J. KRIEK 141–148

Mice

Leishmania major infections in *Phlebotomus duboscqi* fed on murine models immunized with *L. major* sub-cellular antigens and sandfly gut antigens—P.A. MBATI, C.O. ANJILI, S. ODONGO, P. OGAJA and W. TONU 57–63

Suggested dosage rates of melarsoprol in the treatment of mice experimentally infected with *Trypanosoma brucei gambiense*—P.A. MBATI, K. HIRUMI, N. INOUE, N.H. SITUAKIBANZA and H. HIRUMI 71–74

Milk

Morphological changes associated with the development of the rumino-reticulum in growing lambs fed different rations—G.E. SWAN and H.B. GROENEWALD 105–114

Mites

Oribatid mites (Acari, Oribatida) as intermediate hosts of tapeworms of the Family Anoplocephalidae (Cestoda) and the transmission of *Moniezia expansa* cysticercoids in South Africa—R. SCHUSTER, L. COETZEE and J.F. PUTTERILL 49–55

Molineoidea

Molineus cati n. sp. (Nematoda, Trichostrongylina, Molineoidea), a parasite of feral cats, *Felis catus* Linnaeus, 1758 in South Africa—MARIE-CLAUDE DURETTE-DESSET, J. BOOMKER and F.S. MALAN 173–177

Molineus barbaris, Molineus cati, Molineus vexillarius

Molineus cati n. sp. (Nematoda, Trichostrongylina, Molineoidea), a parasite of feral cats, *Felis catus* Linnaeus, 1758 in South Africa—MARIE-CLAUDE DURETTE-DESSET, J. BOOMKER and F.S. MALAN 173–177

Moniezia benedeni

Parasites of South African wildlife. XVI. Helminths of some antelope species from the Eastern and Western Cape Provinces—J. BOOMKER, I.G. HORAK, R. WATERMEYER and D.G. BOOYSE 31–41

Moniezia expansa

Oribatid mites (Acari, Oribatida) as intermediate hosts of tapeworms of the Family Anoplocephalidae (Cestoda) and the transmission of *Moniezia expansa* cysticercoids in South Africa—R. SCHUSTER, L. COETZEE and J.F. PUTTERILL 49–55

Monofluoroacetate
Investigation of biological samples for monofluoroacetate and *Dichapetalum cymosum* poisoning in southern Africa—P.P. MINNAAR, R.I. McCRINDLE, T.W. NAUDÉ and C.J. BOTHA 27–30

Monogenea
Observations on a lesser-known monogenean, *Udonella myliobati*, from a copepod parasite, *Lepeophtheirus natalensis*, parasitizing the spotted ragged-tooth shark, *Carcharias taurus*, from South African waters—P.A.S. OLIVIER, S.M. DIPPENAAR, L.F. KHALIL and N.M. MOKGALONG 135–140

Mozambique
The distribution of tsetse (Diptera: Glossinidae) and bovine trypanosomosis in the Matutuine District, Maputo Province, Mozambique—I. SIGAUQUE, P. VAN DEN BOSSCHE, M. MOIANA, S. JAMAL and L. NEVES 167–172

Muliercula ngoyensis
Oribatid mites (Acari, Oribatida) as intermediate hosts of tapeworms of the Family Anoplocephalidae (Cestoda) and the transmission of *Moniezia expansa* cysticercoids in South Africa—R. SCHUSTER, L. COETZEE and J.F. PUTTERILL 49–55

Murine models
Leishmania major infections in *Phlebotomus duboscqi* fed on murine models immunized with *L. major* sub-cellular antigens and sandfly gut antigens—P.A. MBATI, C.O. ANJILI, S. ODONGO, P. OGAJA and W. TONU 57–63

Mycobacterium bovis
The rise and fall of tuberculosis in a free-ranging chacma baboon troop in the Kruger National Park—D.F. KEET, N.P.J. KRIEK, R.G. BENGIS, D.G. GROBLER and A. MICHEL 115–122

Mycoplasma
Pasteurella gallinarum: Zimbabwean experience of a versatile pathogen—K. MOHAN, F. DZIVA and D. CHITAURO 301–305

N'Dama cattle
In vitro erythrophagocytosis by cultured macrophages stimulated with extraneous substances and those isolated from the blood, spleen and bone marrow of Boran and N'Dama cattle infected with *Trypanosoma congolense* and *Trypanosoma vivax*—V.O. TAIWO and V.O. ANOSA 273–287

Differential expression of surface membrane antigens on bovine monocytes activated with recombinant cytokines and during *Trypanosoma congolense* infection—V.O. TAIWO, V.O. ANOSA and J.O. OLUWANIYI 289–296

Nematoda
Molineus cati n. sp. (Nematoda, Trichostrongylina, Molineoidea), a parasite of feral cats, *Felis catus* Linnaeus, 1758 in South Africa—MARIE-CLAUDE DURETTE-DESSET, J. BOOMKER and F.S. MALAN 173–177

Nematode parasites
A check-list of the nematode parasites of South African serpentes (snakes) and sauria (lizards)—S.F.B.N. HERINGHAGENBECK and J. BOOMKER 1–13

Neodiplozoon polycotyleus
Occurrence of *Neodiplozoon polycotyleus* Paperna, 1973 (Diplozoidae: Monogenea) in cyprinid fish in South Africa—S.N. MASHEGO 153–154

Nile crocodile
Selected chemical parameters in the blood and metals in the organs of the Nile crocodile, *Crocodylus niloticus*, in the Kruger National Park—D. SWANEPOEL, J. BOOMKER and N.P.J. KRIEK 141–148

Ophidascaris
A check-list of the nematode parasites of South African serpentes (snakes) and sauria (lizards)—S.F.B.N. HERINGHAGENBECK and J. BOOMKER 1–13

Oribatida
Oribatid mites (Acari, Oribatida) as intermediate hosts of tapeworms of the Family Anoplocephalidae (Cestoda) and the transmission of *Moniezia expansa* cysticercoids in South Africa—R. SCHUSTER, L. COETZEE and J.F. PUTTERILL 49–55

Osteolaemus tetraspis
Aerobic intestinal flora of wild-caught African dwarf crocodiles *Osteolaemus tetraspis*—F.W. HUCHZERMEYER, M.M. HENTON, J. RILEY and M. AGNAGNA 201–204

Papio ursinus
The rise and fall of tuberculosis in a free-ranging chacma baboon troop in the Kruger National Park—D.F. KEET, N.P.J. KRIEK, R.G. BENGIS, D.G. GROBLER and A. MICHEL 115–122

Parasites

A check-list of the nematode parasites of South African serpentes (snakes) and sauria (lizards)—S.F.B.N. HERINGHAGENBECK and J. BOOMKER 1–13

Parasites of South African wildlife. XVI. Helminths of some antelope species from the Eastern and Western Cape Provinces—J. BOOMKER, I.G. HORAK, R. WATERMEYER and D.G. BOOYSE 31–41

Oribatid mites (Acari, Oribatida) as intermediate hosts of tapeworms of the Family Anoplocephalidae (Cestoda) and the transmission of *Moniezia expansa* cysticercoids in South Africa—R. SCHUSTER, L. COETZEE and J.F. PUTTERILL 49–55

Helminth parasites of domestic pigeons (*Columba livia domestica*) in Sebele, Gaborone, Botswana—E.Z. MUSHI, M.G. BINTA, R.G. CHABO, R. NDEBELE and R. PANZIRAH 75–76

Observations on a lesser-known monogenean, *Udonella myliobati*, from a copepod parasite, *Lepeophtheirus natalensis*, parasitizing the spotted ragged-tooth shark, *Carcharias taurus*, from South African waters—P.A.S. OLIVIER, S.M. DIPPENAAR, L.F. KHALIL and N.M. MOKGALONG 135–140

Occurrence of *Neodiplozoon polycotyleus* Paperna, 1973 (Diplozoidae: Monogenea) in cyprinid fish in South Africa—S.N. MASHEGO 153–154

The role of parasitic diseases as causes of mortality in cattle in a high potential area of central Kenya: a quantitative analysis—P.W.N. KANYARI and J. KAGIRA 157–161

Molineus cati n. sp. (Nematoda, Trichostrongylina, Molineoidea), a parasite of feral cats, *Felis catus* Linnaeus, 1758 in South Africa—MARIE-CLAUDE DURETTE-DESSET, J. BOOMKER and F.S. MALAN 173–177

The prevalence of internal and external parasites in pigs of different ages and sexes in Southeast District, Botswana—S.J. NSOSO, K.P. MOSALA, R.T. NDEBELE and S.S. RAMABU 217–220

Studies on the genus *Setaria* Viborg, 1795 in South Africa. I. *Setaria africana* (Yeh, 1959)—R. WATERMEYER, J. BOOMKER and J.F. PUTTERILL 229–234

Parasites of domestic and wild animals in South Africa. XXXVIII. Ixodid ticks collected from 23 wild carnivore species—I.G. HORAK, L.E.O. BRAACK, L.J. FOURIE and JANE B. WALKER 239–250

Pasteurella multocida

Capsular serogroups of *Pasteurella multocida* isolated from animals in Zimbabwe—F. DZIVA, K. MOHAN and A. PAWANDIWA 225–228

PCR

Suggested dosage rates of melarsoprol in the treatment of mice experimentally infected with *Trypanosoma brucei gambiense*—P.A. MBATI, K. HIRUMI, N. INOUE, N.H. SITUAKIBANZA and H. HIRUMI 71–74

Phage types

Phage types of *Salmonella enterica* serovar Enteritidis isolated in South Africa from 1991–1995—L. MARÉ, M.L. VAN DER WALT and L.M.T. DICKS 129–133

Phlebotomus duboscqi

Leishmania major infections in *Phlebotomus duboscqi* fed on murine models immunized with *L. major* sub-cellular antigens and sandfly gut antigens—P.A. MBATI, C.O. ANJILI, S. ODONGO, P. OGAJA and W. TONU 57–63

Pigeons

Helminth parasites of domestic pigeons (*Columba livia domestica*) in Sebele, Gaborone, Botswana—E.Z. MUSHI, M.G. BINTA, R.G. CHABO, R. NDEBELE and R. PANZIRAH 75–76

Pigs

Phage types of *Salmonella enterica* serovar Enteritidis isolated in South Africa from 1991–1995—L. MARÉ, M.L. VAN DER WALT and L.M.T. DICKS 129–133

The prevalence of internal and external parasites in pigs of different ages and sexes in Southeast District, Botswana—S.J. NSOSO, K.P. MOSALA, R.T. NDEBELE and S.S. RAMABU 217–220

Poison

Investigation of biological samples for monofluoroacetate and *Dichapetalum cymosum* poisoning in southern Africa—P.P. MINNAAR, R.I. McCRINDLE, T.W. NAUDÉ and C.J. BOTHA 27–30

Poultry

Phage types of *Salmonella enterica* serovar Enteritidis isolated in South Africa from 1991–1995—L. MARÉ, M.L. VAN DER WALT and L.M.T. DICKS 129–133

Prunus africana

Biochemical and haematological changes in rats administered an aqueous extract of *Prunus africana* stem bark at various dosage levels—P.K. GATHUMBI, J.W. MWANGI, S.M. NJIRO and G.M. MUGERA 123–128

Pygeum africanum
Biochemical and haematological changes in rats administered an aqueous extract of *Prunus africana* stem bark at various dosage levels—P.K. GATHUMBI, J.W. MWANGI, S.M. NJIRO and G.M. MUGERA 123–128

Rafoxanide
Stability and partitioning of closantel and rafoxanide in ruminal fluid of sheep—G.E. SWAN, D.W. OLIVER, J. VAN RENSBURG, H.S. STEYN and MARIA S.G. MÜLDERS 97–103

***Raillietina* spp.**
Helminth parasites of domestic pigeons (*Columba livia domestica*) in Sebele, Gaborone, Botswana—E.Z. MUSHI, M.G. BINTA, R.G. CHABO, R. NDEBELE and R. PANZIRAH 75–76

Rats
Biochemical and haematological changes in rats administered an aqueous extract of *Prunus africana* stem bark at various dosage levels—P.K. GATHUMBI, J.W. MWANGI, S.M. NJIRO and G.M. MUGERA 123–128

Fertility in rats immunized with steroid-free bovine follicular fluid—C.R. SEWANI-RUSIKE and C. DAKWA 257–262

Redunca fulvorufula
Parasites of South African wildlife. XVI. Helminths of some antelope species from the Eastern and Western Cape Provinces—J. BOOMKER, I.G. HORAK, R. WATERMEYER and D.G. BOOYSE 31–41

Reptiles
A check-list of the nematode parasites of South African serpentes (snakes) and sauria (lizards)—S.F.B.N. HERINGHAGENBECK and J. BOOMKER 1–13

RT-PCR
The prevalence of different African horsesickness virus serotypes in the Onderstepoort area near Pretoria, during an outbreak of African horsesickness in South Africa in 1995/1996—C.W. BREMER, G.H. GERDES, H. AITCHISON, I. LOUW, R.R. GREYLING and J. WELGEMOED 65–70

Rumen, rumino-reticulum
Morphological changes associated with the development of the rumino-reticulum in growing lambs fed different rations—G.E. SWAN and H.B. GROENEWALD 105–114

Ruminal fluid
Stability and partitioning of closantel and rafoxanide in ruminal fluid of sheep—G.E. SWAN, D.W. OLIVER, J. VAN RENSBURG, H.S. STEYN and MARIA S.G. MÜLDERS 97–103

Salicylanilides
Stability and partitioning of closantel and rafoxanide in ruminal fluid of sheep—G.E. SWAN, D.W. OLIVER, J. VAN RENSBURG, H.S. STEYN and MARIA S.G. MÜLDERS 97–103

Salmonella
Aerobic intestinal flora of wild-caught African dwarf crocodiles Osteolaemus tetraspis—F.W. HUCHZERMEYER, M.M. HENTON, J. RILEY and M. AGNAGNA 201–204

Salmonella enterica
Phage types of *Salmonella enterica* serovar Enteritidis isolated in South Africa from 1991–1995—L. MARÉ, M.L. VAN DER WALT and L.M.T. DICKS 129–133

Salmonella gallinarum
Bacterial colonization and endotoxin activity during experimental acute fowl typhoid in chickens—T. KOKOSHAROV 297–300

Sandfly
Leishmania major infections in *Phlebotomus duboscqi* fed on murine models immunized with *L. major* sub-cellular antigens and sandfly gut antigens—P.A. MBATI, C.O. ANJILI, S. ODONGO, P. OGAJA and W. TONU 57–63

Sarcoptes scabiei
The prevalence of internal and external parasites in pigs of different ages and sexes in Southeast District, Botswana—S.J. NSOSO, K.P. MOSALA, R.T. NDEBELE and S.S. RAMABU 217–220

Sauria
A check-list of the nematode parasites of South African serpentes (snakes) and sauria (lizards)—S.F.B.N. HERINGHAGENBECK and J. BOOMKER 1–13

Scheloribates fusifer
Oribatid mites (Acari, Oribatida) as intermediate hosts of tapeworms of the Family Anoplocephalidae (Cestoda and the transmission of *Moniezia expansa* cysticercoids in South Africa—R. SCHUSTER, L. COETZEE and J.F. PUTTERILL 49–55

SEM
Ultrastructural study of the luminal surface of the ducts of the epididymis of gallinaceous birds—T.A. AIRE and D. JOSLING 191–199

Serology
Evaluation of a commercially available ELISA kit for detection of antibodies to *Anaplasma marginale* and *Anaplasma centrale* in cattle in Australia and Zimbabwe—P.M. BOWLES, J.B. MOLLOY, G.W. BLIGHT, S. SINGH and L.G. MABIKACHECHE 83–86

Serotypes
The prevalence of different African horsesickness virus serotypes in the Onderstepoort area near Pretoria, during an outbreak of African horsesickness in South Africa in 1995/1996—C.W. BREMER, G.H. GERDES, H. AITCHISON, I. LOUW, R.R. GREYLING and J. WELGEMOED 65–70

Serpentes
A check-list of the nematode parasites of South African serpentes (snakes) and sauria (lizards)—S.F.B.N. HERINGHAGENBECK and J. BOOMKER 1–13

Setaria africana
Studies on the genus Setaria Viborg, 1795 in South Africa. I. *Setaria africana* (Yeh, 1959)—R. WATERMEYER, J. BOOMKER and J.F. PUTTERILL 229–234

Shark
Observations on a lesser-known monogenean, *Udonella myliobati*, from a copepod parasite, *Lepeophtheirus natalensis*, parasitizing the spotted ragged-tooth shark, *Carcharias taurus*, from South African waters—P.A.S. OLIVIER, S.M. DIPPENAAR, L.F. KHALIL and N.M. MOKGALONG 135–1400

Sheep
Suspected sheep-associated malignant catarrhal fever in a zero-grazed dairy herd in Kenya—C.M MULEI, P.K. GATHUMBI and P.G. MBUTHIA 43–47

The effect of a natural maedi-visna virus infection on the productivity of South African sheep—B. DUNGU, J. VORSTER, G.F. BATH and D.W. VERWOERD 87–96

Stability and partitioning of closantel and rafoxanide in ruminal fluid of sheep—G.E. SWAN, D.W. OLIVER, J. VAN RENSBURG, H.S. STEYN and MARIA S.G. MÜLDERS 97–103

Phage types of *Salmonella enterica* serovar Enteritidis isolated in South Africa from 1991–1995—L. MARÉ, M.L. VAN DER WALT and L.M.T. DICKS 129–133

Snakes
A check-list of the nematode parasites of South African serpentes (snakes) and sauria (lizards)—S.F.B.N. HERINGHAGENBECK and J. BOOMKER 1–13

South Africa
A check-list of the nematode parasites of South African serpentes (snakes) and sauria (lizards)—S.F.B.N. HERINGHAGENBECK and J. BOOMKER 1–13

A newly developed odour-baited "H trap" for the live collection of *Glossina brevipalpis* and *Glossina austeni* (Diptera: Glossinidae) in South Africa—KARIN KAPPMEIER 15–26

Investigation of biological samples for monofluoroacetate and *Dichapetalum cymosum* poisoning in southern Africa—P.P. MINNAAR, R.I. MCCRINDLE, T.W. NAUDÉ and C.J. BOTHA 27–30

Parasites of South African wildlife. XVI. Helminths of some antelope species from the Eastern and Western Cape Provinces—J. BOOMKER, I.G. HORAK, R. WATERMEYER and D.G. BOOYSE 31–41

Oribatid mites (Acari, Oribatida) as intermediate hosts of tapeworms of the Family Anoplocephalidae (Cestoda and the transmission of *Moniezia expansa* cysticercoids in South Africa—R. SCHUSTER, L. COETZEE and J.F. PUTTERILL 49–55

The prevalence of different African horsesickness virus serotypes in the Onderstepoort area near Pretoria, during an outbreak of African horsesickness in South Africa in 1995/1996—C.W. BREMER, G.H. GERDES, H. AITCHISON, I. LOUW, R.R. GREYLING and J. WELGEMOED 65–70

The effect of a natural maedi-visna virus infection on the productivity of South African sheep—B. DUNGU, J. VORSTER, G.F. BATH and D.W. VERWOERD 87–96

The rise and fall of tuberculosis in a free-ranging chacma baboon troop in the Kruger National Park—D.F. KEET, N.P.J. KRIEK, R.G. BENGIS, D.G. GROBLER and A. MICHEL 115–122

Phage types of *Salmonella enterica* serovar Enteritidis isolated in South Africa from 1991–1995—L. MARÉ, M.L. VAN DER WALT and L.M.T. DICKS 129–133

Observations on a lesser-known monogenean, *Udonella myliobati*, from a copepod parasite, *Lepeophtheirus natalensis*, parasitizing the spotted ragged-tooth shark, *Carcharias taurus*, from South African waters—P.A.S. OLIVIER, S.M. DIPPENAAR, L.F. KHALIL and N.M. MOKGALONG 135–140

Occurrence of *Neodiplozoon polycotyleus* Paperna, 1973 (Diplozoidae: Monogenea) in cyprinid fish in South Africa—S.N. MASHEGO 153–154

Molineus cati n. sp. (Nematoda, Trichostrongylina, Molineoidea), a parasite of feral cats, *Felis catus* Linnaeus, 1758 in South Africa—MARIE-CLAUDE DURETTE-DESSET, J. BOOMKER and F.S. MALAN 173–177

Diurnal activity patterns of *Glossina brevipalpis* and *G. austeni* (Diptera: Glossinidae) in South Africa, with reference to season and meteorological factors—KARIN KAPPMEIER 179–189

Studies on the genus *Setaria* Viborg, 1795 in South Africa. I. *Setaria africana* (Yeh, 1959)—R. WATERMEYER, J. BOOMKER and J.F. PUTTERILL 229–234

Parasites of domestic and wild animals in South Africa. XXXVIII. Ixodid ticks collected from 23 wild carnivore species—I.G. HORAK, L.E.O. BRAACK, L.J. FOURIE and JANE B. WALKER 239–250

Parameters for the determination and evaluation of heat stress in dairy cattle in South Africa—J.H. DU PREEZ 263–271

Sudan

Application of ultrasonography in prevalence studies of hydatid cysts in goats in north-western Turkana, Kenya and Toposaland, southern Sudan—E.M. NJOROGE, P.M.F. MBITHI, J.M. GATHUMA, T.M. WACHIRA, J.K. MAGAMBO and E. ZEYHLE 251–255

Spauligodon morgani, Spauligodon vojteki

A check-list of the nematode parasites of South African serpentes (snakes) and sauria (lizards)—S.F.B.N. HERING-HAGENBECK and J. BOOMKER 1–13

Stem-bark

Biochemical and haematological changes in rats administered an aqueous extract of *Prunus africana* stem bark at various dosage levels—P.K. GATHUMBI, J.W. MWANGI, S.M. NJIRO and G.M. MUGERA 123–128

Syncerus caffer

The rise and fall of tuberculosis in a free-ranging chacma baboon troop in the Kruger National Park—D.F. KEET, N.P.J. KRIEK, R.G. BENGIS, D.G. GROBLER and A. MICHEL 115–122

Tamarinus nigrocollis

Molineus cati n. sp. (Nematoda, Trichostrongylina, Molineoidea), a parasite of feral cats, *Felis catus* Linnaeus, 1758 in South Africa—MARIE-CLAUDE DURETTE-DESSET, J. BOOMKER and F.S. MALAN 173–177

Tapeworms

Oribatid mites (Acari, Oribatida) as intermediate hosts of tapeworms of the Family Anoplocephalidae (Cestoda and the transmission of *Moniezia expansa* cysticercoids in South Africa—R. SCHUSTER, L. COETZEE and J.F. PUTTERILL49–55

Taurotragus oryx

Parasites of South African wildlife. XVI. Helminths of some antelope species from the Eastern and Western Cape Provinces—J. BOOMKER, I.G. HORAK, R. WATERMEYER and D.G. BOOYSE 31–41

Tayra barbara

Molineus cati n. sp. (Nematoda, Trichostrongylina, Molineoidea), a parasite of feral cats, *Felis catus* Linnaeus, 1758 in South Africa—MARIE-CLAUDE DURETTE-DESSET, J. BOOMKER and F.S. MALAN 173–177

TEM

Ultrastructural study of the luminal surface of the ducts of the epididymis of gallinaceous birds—T.A. AIRE and D. JOSLING 191–199

Theileria mutans

Bovine borreliosis in Botswana—S.P. SHARMA, W. AMANFU and T.C. LOSHO 221–223

Tick-borne diseases

The role of parasitic diseases as causes of mortality in cattle in a high potential area of central Kenya: a quantitative analysis—P.W.N. KANYARI and J. KAGIRA 157–161

Ticks

Efficacy of flumethrin 1 % pour-on against ticks on cattle under field condition in Ethiopia—S. MEKONNEN 235–237

Parasites of domestic and wild animals in South Africa. XXXVIII. Ixodid ticks collected from 23 wild carnivore species—I.G. HORAK, L.E.O. BRAACK, L.J. FOURIE and JANE B. WALKER 239–250

Toposaland

Application of ultrasonography in prevalence studies of hydatid cysts in goats in north-western Turkana, Kenya and Toposaland, southern Sudan—E.M. NJOROGE, P.M.F. MBITHI, J.M. GATHUMA, T.M. WACHIRA, J.K. MAGAMBO and E. ZEYHLE 251–255

Trap

A newly developed odour-baited "H trap" for the live collection of *Glossina brevipalpis* and *Glossina austeni* (Diptera: Glossinidae) in South Africa—KARIN KAPPMEIER 15–26

Trichostrongylina

Molineus cati n. sp. (Nematoda, Trichostrongylina, Molineoidea), a parasite of feral cats, *Felis catus* Linnaeus, 1758 in South Africa—MARIE-CLAUDE DURETTE-DESSET, J. BOOMKER and F.S. MALAN 173–177

***Trichostrongylus* spp.**

The prevalence of internal and external parasites in pigs of different ages and sexes in Southeast District, Botswana—S.J. NSOSO, K.P. MOSALA, R.T. NDEBELE and S.S. RAMABU 217–220

Trichuris suis

The prevalence of internal and external parasites in pigs of different ages and sexes in Southeast District, Botswana—S.J. NSOSO, K.P. MOSALA, R.T. NDEBELE and S.S. RAMABU 217–220

Trypanosoma

T. brucei gambiense

Suggested dosage rates of melarsoprol in the treatment of mice experimentally infected with *Trypanosoma brucei gambiense*—P.A. MBATI, K. HIRUMI, N. INOUE, N.H. SITUAKIBANZA and H. HIRUMI 71–74

T. congolense

The distribution of tsetse (Diptera: Glossinidae) and bovine trypanosomosis in the Matutuine District, Maputo Province, Mozambique—I. SIGAUQUE, P. VAN DEN BOSSCHE, M. MOIANA, S. JAMAL and L. NEVES 167–172

In vitro erythrophagocytosis by cultured macrophages stimulated with extraneous substances and those isolated from the blood, spleen and bone marrow of Boran and N'Dama cattle infected with *Trypanosoma congolense* and *Trypanosoma vivax*—V.O. TAIWO and V.O. ANOSA 273–287

T. vivax

In vitro erythrophagocytosis by cultured macrophages stimulated with extraneous substances and those isolated from the blood, spleen and bone marrow of Boran and N'Dama cattle infected with *Trypanosoma congolense* and *Trypanosoma vivax*—V.O. TAIWO and V.O. ANOSA 273–287

Trypanosomosis

The distribution of tsetse (Diptera: Glossinidae) and bovine trypanosomosis in the Matutuine District, Maputo Province, Mozambique—I. SIGAUQUE, P. VAN DEN BOSSCHE, M. MOIANA, S. JAMAL and L. NEVES 167–172

Trypanotolerance

Differential expression of surface membrane antigens on bovine monocytes activated with recombinant cytokines and during *Trypanosoma congolense* infection—V.O. TAIWO, V.O. ANOSA and J.O. OLUWANIY 289–296

Tsetse

A newly developed odour-baited "H trap" for the live collection of *Glossina brevipalpis* and *Glossina austeni* (Diptera: Glossinidae) in South Africa—KARIN KAPPMEIER 15–26

The distribution of tsetse (Diptera: Glossinidae) and bovine trypanosomosis in the Matutuine District, Maputo Province, Mozambique—I. SIGAUQUE, P. VAN DEN BOSSCHE, M. MOIANA, S. JAMAL and L. NEVES 167–172

Diurnal activity patterns of *Glossina brevipalpis* and *G. austeni* (Diptera: Glossinidae) in South Africa, with reference to season and meteorological factors—KARIN KAPPMEIER 179–189

Tuberculosis

The rise and fall of tuberculosis in a free-ranging chacma baboon troop in the Kruger National Park—D.F. KEET, N.P.J. KRIEK, R.G. BENGIS, D.G. GROBLER and A. MICHEL 115–122

Turkana

Application of ultrasonography in prevalence studies of hydatid cysts in goats in north-western Turkana, Kenya and Toposaland, southern Sudan—E.M. NJOROGE, P.M.F. MBITHI, J.M. GATHUMA, T.M. WACHIRA, J.K. MAGAMBO and E. ZEYHLE 251–255

Udonella myliobati

Observations on a lesser-known monogenean, *Udonella myliobati*, from a copepod parasite, *Lepeophtheirus natalensis*, parasitizing the spotted ragged-tooth shark, *Carcharias taurus*, from South African waters—P.A.S. OLIVIER, S.M. DIPPENAAR, L.F. KHALIL and N.M. MOKGALONG 135–140

Vanillic acid

Parameters for the determination and evaluation of heat stress in dairy cattle in South Africa—J.H. DU PREEZ 263–271

Varanus niloticus

A check-list of the nematode parasites of South African serpentes (snakes) and sauria (lizards)—S.F.B.N. HERINGHAGENBECK and J. BOOMKER 1–13

Virus

The prevalence of different African horsesickness virus serotypes in the Onderstepoort area near Pretoria, during an outbreak of African horsesickness in South Africa in 1995/1996—C.W. BREMER, G.H. GERDES, H. AITCHISON, I. LOUW, R.R. GREYLING and J. WELGEMOED 65–70

The effect of a natural maedi-visna virus infection on the productivity of South African sheep—B. DUNGU, J. VORSTER, G.F. BATH and D.W. VERWOERD 87–96

Wildlife

Parasites of South African wildlife. XVI. Helminths of some antelope species from the Eastern and Western Cape Provinces—J. BOOMKER, I.G. HORAK, R. WATERMEYER and D.G. BOOYSE 31–41

Studies on the genus *Setaria* Viborg, 1795 in South Africa. I. *Setaria africana* (Yeh, 1959)—R. WATERMEYER, J. BOOMKER and J.F. PUTTERILL 229–234

Parasites of domestic and wild animals in South Africa. XXXVIII. Ixodid ticks collected from 23 wild carnivore species—I.G. HORAK, L.E.O. BRAACK, L.J. FOURIE and JANE B. WALKER 239–250

Worms

Observations on a lesser-known monogenean, *Udonella myliobati*, from a copepod parasite, *Lepeophtheirus natalensis*, parasitizing the spotted ragged-tooth shark, *Carcharias taurus*, from South African waters—P.A.S. OLIVIER, S.M. DIPPENAAR, L.F. KHALIL and N.M. MOKGALONG 135–140

Zebra

The prevalence of different African horsesickness virus serotypes in the Onderstepoort area near Pretoria, during an outbreak of African horsesickness in South Africa in 1995/1996—C.W. BREMER, G.H. GERDES, H. AITCHISON, I. LOUW, R.R. GREYLING and J. WELGEMOED 65–70

Zero grazed

Suspected sheep-associated malignant catarrhal fever in a zero-grazed dairy herd in Kenya—C.M MULEI, P.K. GATHUMBI and P.G. MBUTHIA 43–47

Zimbabwe

Evaluation of a commercially available ELISA kit for detection of antibodies to *Anaplasma marginale* and *Anaplasma centrale* in cattle in Australia and Zimbabwe—P.M. BOWLES, J.B. MOLLOY, G.W. BLIGHT, S. SINGH and L.G. MABIKACHECHE 83–86

Capsular serogroups of *Pasteurella multocida* isolated from animals in Zimbabwe—F. DZIVA, K. MOHAN and A. PAWANDIWA 225–228